

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number  
**WO 01/12604 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 213/61**,  
213/89, 405/12, 213/77, 401/12, 213/81, 213/64, 409/12,  
417/12, 498/04, 401/06, A01N 43/40

(GB). **PATEL, Gita** [GB/GB]; Chesterfield House, Wulfstan Way, Cambridge CB1 8QL (GB). **SCHNATTERER, Stefan** [DE/DE]; Schillerring 10, D-65795 Hattersheim (DE).

(21) International Application Number: **PCT/EP00/08268**

(22) International Filing Date: 11 August 2000 (11.08.2000)

(74) Agent: **MERIGEAULT, Shona**; Aventis CropScience S.A., B.P. 9163, F-69263 Lyon Cedex 09 (FR).

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:  
9919558.8 18 August 1999 (18.08.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **AVENTIS CROPSCIENCE GMBH** [DE/DE]; Brünningstrasse 50, D-65929 Frankfurt am Main (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **COOKE, Tracey** [GB/GB]; 7 Larch Avenue, Brickett Wood, St Albans AL2 3SN (GB). **HARDY, David** [GB/GB]; 46 St. Bedes Gardens, Cambridge CB1 3UF (GB). **MOLONEY, Brian, Anthony** [GB/GB]; 2 Crookdale Beck, Oxon OX11 7US (GB). **O'MAHONY, Mary, Josephine** [GB/GB]; 8 Manger's Lane, Duxford, Cambridge CB2 4RN (GB). **PETTETT, Michael, George** [GB/GB]; Aventis CropScience UK Ltd., Fyfield Road, Ongar Essex CM5 OHW

**Published:**

— *With international search report.*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **FUNGICIDES**



(I)


(57) Abstract: Use of compounds of general formula (I) or salts thereof as phytopathogenic fungicides wherein the various radicals and substituents are as defined in the description, pesticidal compositions containing them and method for combatting pests which comprises applying these.

WO 01/12604 A1

Fungicides

5 This invention relates to compounds having fungicidal activity.

In a first aspect the invention provides the use of a compound of general formula I or salts thereof as phytopathogenic fungicides



10

where

$A^1$  is 2-pyridyl or its *N*-oxide, each of which may be substituted by up to four groups at least one of which is haloalkyl;

15  $A^2$  is optionally substituted heterocyclyl or optionally substituted carbocyclyl ( $A^2$  is preferably phenyl, cyclohexyl, cyclopropyl or heterocyclyl, each of which may be substituted);

20  $L$  is a 3-atom linker selected from the list:  $-\text{CH}(\text{R}^1)\text{N}(\text{R}^3)\text{CH}(\text{R}^2)-$ ,  $-\text{N}(\text{R}^3)\text{N}(\text{R}^4)\text{C}(=\text{X})-$ ,  $-\text{C}(=\text{X})\text{N}(\text{R}^3)\text{CH}(\text{R}^1)-$ ,  $-\text{CH}(\text{R}^1)\text{OC}(=\text{X})-$ ,  $-\text{CH}(\text{R}^1)\text{OCH}(\text{R}^2)-$ ,  $-\text{N}(\text{R}^3)\text{C}(=\text{X})\text{N}(\text{R}^4)-$ ,  $-\text{C}(\text{R}^1)=\text{C}(\text{R}^2)\text{C}(=\text{X})-$ ,  $-\text{C}(\text{R}^1)=\text{N}-\text{N}(\text{R}^3)-$ ,  $-\text{CH}(\text{R}^1)\text{N}=\text{C}(\text{R}^2)-$ ,  $-\text{O}-\text{N}=\text{C}(\text{R}^1)-$ ,  $-\text{O}-\text{N}(\text{R}^3)\text{C}(=\text{X})-$ ,  $-\text{N}(\text{R}^3)\text{N}(\text{R}^4)\text{CH}(\text{R}^1)-$ ,  $-\text{N}(\text{R}^3)\text{C}(\text{Y})=\text{N}-$ ,  $-\text{N}=\text{C}(\text{Y})-\text{N}(\text{R}^3)-$ ,  $-\text{N}(\text{R}^3)\text{N}=\text{C}(\text{Y})-$ ,  $-\text{C}(=\text{X})-\text{N}(\text{R}^3)\text{N}(\text{R}^4)-$ ,  $-\text{C}(\text{Y})=\text{N}-\text{N}(\text{R}^4)-$  and  $-\text{N}(\text{R}^3)\text{CH}(\text{R}^1)\text{C}(=\text{X})-$ ; wherein  $A^1$  is attached to the left hand side of linker  $L$  ( $L$  is preferably selected from the list:  $-\text{CH}(\text{R}^1)\text{N}(\text{R}^3)\text{CH}(\text{R}^2)-$ ,  $-\text{N}(\text{R}^3)\text{N}(\text{R}^4)\text{C}(=\text{X})-$ ,  $-\text{C}(=\text{X})\text{N}(\text{R}^3)\text{CH}(\text{R}^1)-$ ,  $-\text{CH}(\text{R}^1)\text{OC}(=\text{X})-$ ,  $-\text{CH}(\text{R}^1)\text{OCH}(\text{R}^2)-$ ,  $-\text{N}(\text{R}^3)\text{C}(=\text{X})\text{N}(\text{R}^4)-$ ,  $-\text{C}(\text{R}^1)=\text{C}(\text{R}^2)\text{C}(=\text{X})-$ ,  $-\text{C}(\text{R}^1)=\text{N}-\text{N}(\text{R}^3)-$ ,  $-\text{CH}(\text{R}^1)\text{N}=\text{C}(\text{R}^2)-$ ,  $-\text{O}-\text{N}=\text{C}(\text{R}^1)-$ ,  $-\text{O}-\text{N}(\text{R}^3)\text{C}(=\text{X})-$ );

25

where  $\text{R}^1$  and  $\text{R}^2$ , which may be the same or different, are  $\text{R}^b$ , cyano, nitro, halogen,  $-\text{OR}^b$ ,  $-\text{SR}^b$  or optionally substituted amino ( $\text{R}^1$  and  $\text{R}^2$  are preferably hydrogen, acyl, optionally substituted alkyl, cyano or optionally substituted phenyl);

$R^3$  and  $R^4$ , which may be the same or different, are  $R^b$ , cyano or nitro ( $R^3$  and  $R^4$  are preferably hydrogen, acyl or optionally substituted alkyl);

or any  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  group, together with the interconnecting atoms, can form a 5- or 6-membered ring with any other  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$ , or any  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  group, together with the interconnecting atoms can form a 5- or 6-membered ring with  $A^2$ ;

X is oxygen, sulfur,  $N-OR^b$ ,  $N-R^b$  or  $N-N(R^b)_2$  (X is preferably oxygen or sulfur); and

Y is halogen,  $-OR^b$ ,  $-SR^b$ ,  $-N(R^b)_2$ ,  $-NR^b(OR^b)$  or  $-NR^bN(R^b)_2$ ;

wherein  $R^b$  is alkyl, alkenyl, alkynyl, carbocyclyl or heterocyclyl, each of which may be substituted; or hydrogen or acyl, or two adjacent  $R^b$  groups together with the nitrogen atom to which they are attached may form a 5- or 6-membered ring.

Preferred substituents on the 2-pyridyl group ( $A^1$ ) are halogen, hydroxy, cyano, nitro,  $SF_5$ , trialkylsilyl, optionally substituted amino, acyl, or a group  $-R^a$ ,  $-OR^a$  or  $-SR^a$ , or a group  $-C(R^a)=N-Q$ , where Q is  $-R^a$ ,  $-OR^a$ ,  $-SR^a$  or optionally substituted amino, wherein  $R^a$  is alkyl, alkenyl, alkynyl, carbocyclyl or heterocyclyl, each of which may be substituted; or two adjacent substituents together with the atoms to which they are attached form an optionally substituted ring which can contain up to 3 hetero atoms. Especially preferred substituents are alkoxy, alkyl, cyano, halogen, nitro, alkoxycarbonyl, alkylsulfinyl, alkylsulfonyl and trifluoromethyl, particularly chlorine and trifluoromethyl.

Preferably, the 2-pyridyl group is substituted at the 3 and/or 5 position.

The invention also includes any of the compounds specifically exemplified hereinafter.

Any alkyl group may be straight or branched and is preferably of 1 to 10 carbon atoms, especially 1 to 7 and particularly 1 to 5 carbon atoms.

Any alkenyl or alkynyl group may be straight or branched and is preferably of 2 to 7 carbon atoms and may contain up to 3 double or triple bonds which may be conjugated, for example vinyl, allyl, butadienyl or propargyl.

Any carbocyclyl group may be saturated, unsaturated or aromatic, and contain 3 to 8 ring-atoms. Preferred saturated carbocyclyl groups are cyclopropyl, cyclopentyl or cyclohexyl. Preferred unsaturated carbocyclyl groups contain up to 3 double bonds. A preferred aromatic carbocyclyl group is phenyl. The term carbocyclic should be similarly construed. In addition, the term carbocyclyl includes any fused combination of carbocyclyl groups, for example naphthyl, phenanthryl, indanyl and indenyl.

Any heterocyclyl group may be saturated, unsaturated or aromatic, and contain 5 to 7 ring-atoms up to 4 of which may be hetero-atoms such as nitrogen, oxygen and sulfur. Examples of heterocyclyl groups are furyl, thienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, dioxolanyl, oxazolyl, thiazolyl, imidazolyl, imidazoliny, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyranly, pyridyl, piperidinyl, dioxanyl, morpholino, dithianyl, thiomorpholino, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, sulfolanyl, tetrazolyl, triazinyl, azepinyl, oxazepinyl, thiazepinyl, diazepinyl and thiazolinyl. In addition, the term heterocyclyl includes fused heterocyclyl groups, for example benzimidazolyl, benzoxazolyl, imidazopyridinyl, benzoxazinyl, benzothiazinyl, oxazolopyridinyl, benzofuranyl, quinolinyl, quinazolinyl, quinoxalinyl, dihydroquinazolinyl, benzothiazolyl, phthalimido, benzofuranyl, benzodiazepinyl, indolyl and isoindolyl. The term heterocyclic should be similarly construed.

Any alkyl, alkenyl, alkynyl, carbocyclyl or heterocyclyl group, when substituted, may be substituted by one or more substituents, which may be the same or different, and may be selected from the list: hydroxy; mercapto; azido; nitro; halogen; cyano; acyl; optionally substituted amino; optionally substituted carbocyclyl; optionally substituted heterocyclyl; cyanato; thiocyanato;  $-SF_5$ ;  $-OR^a$ ;  $-SR^a$  and  $-Si(R^a)_3$ , where  $R^a$  is alkyl, alkenyl, alkynyl, carbocyclyl or heterocyclyl, each of which may be substituted. In the case of any carbocyclyl or heterocyclyl group the list includes additionally: alkyl, alkenyl and alkynyl, each of which may be substituted. Preferred substituents on any alkyl, alkenyl or alkynyl group are alkoxy, haloalkoxy or alkylthio, each containing 1 to 5 carbon atoms; halogen; or optionally substituted phenyl. Preferred substituents on any carbocyclyl or heterocyclyl

group are alkyl, haloalkyl, alkoxy, haloalkoxy or alkylthio, each containing 1 to 5 carbon atoms; halogen; or optionally substituted phenyl.

5 In the case of any alkyl group or any unsaturated ring-carbon in any carbocyclyl or heterocyclyl group the list includes a divalent group such as oxo or imino, which may be substituted by optionally substituted amino,  $R^a$  or  $-OR^a$ . Preferred groups are oxo, imino, alkylimino, oximino, alkyloximino or hydrazono.

10 Any amino group, when substituted and where appropriate, may be substituted by one or two substituents which may be the same or different, selected from the list: optionally substituted alkyl, optionally substituted amino,  $-OR^a$  and acyl groups. Alternatively two substituents together with the nitrogen to which they are attached may form a heterocyclyl group, preferably a 5 to 7-membered heterocyclyl group, which may be substituted and may contain other hetero atoms, for example morpholino, thiomorpholino or piperidinyl.

15 The term acyl includes the residues of sulfur and phosphorus-containing acids as well as carboxylic acids. Typically the residues are covered by the general formulae  $-C(=X^a)R^c$ ,  $-S(O)_pR^c$  and  $-P(=X^a)(OR^a)(OR^a)$ , where appropriate  $X^a$  is O or S,  $R^c$  is as defined for  $R^a$ ,  $-OR^a$ ,  $-SR^a$ , optionally substituted amino or acyl; and p is 1 or 2. Preferred groups are –  
20  $C(=O)R^d$ ,  $-C(=S)R^d$ , and  $-S(O)_pR^d$  where  $R^d$  is alkyl,  $C_1$  to  $C_5$  alkoxy,  $C_1$  to  $C_5$  alkylthio, phenyl, heterocyclyl or amino, each of which may be substituted.

Complexes of compounds of the invention are usually formed from a salt of formula  $MAN_2$ , in which M is a divalent metal cation, e.g. copper, manganese, cobalt, nickel, iron or zinc  
25 and An is an anion, e.g. chloride, nitrate or sulfate.

In cases where the compounds of the invention exist as the E and Z isomers, the invention includes individual isomers as well as mixtures thereof.

30 In cases where compounds of the invention exist as tautomeric isomers, the invention includes individual tautomers as well as mixtures thereof.

In cases where the compounds of the invention exist as optical isomers, the invention includes individual isomers as well as mixtures thereof.

The compounds of the invention have activity as fungicides, especially against fungal diseases of plants, e.g. mildews and particularly cereal powdery mildew (*Erysiphe graminis*) and vine downy mildew (*Plasmopara viticola*), rice blast (*Pyricularia oryzae*), cereal eyespot (*Pseudocercospora herpotrichoides*), rice sheath blight (*Pellicularia sasakii*), grey mould (*Botrytis cinerea*), damping off (*Rhizoctonia solani*), wheat brown rust (*Puccinia recondita*), late tomato or potato blight (*Phytophthora infestans*), apple scab (*Venturia inaequalis*), and glume blotch (*Leptosphaeria nodorum*). Other fungi against which the compounds may be active include other powdery mildews, other rusts, and other general pathogens of Deuteromycete, Ascomycete, Phycomycete and Basidiomycete origin.

The invention thus also provides a method of combating fungi at a locus infested or liable to be infested therewith, which comprises applying to the locus a compound of formula I.

The invention also provides an agricultural composition comprising a compound of formula I in admixture with an agriculturally acceptable diluent or carrier.

The composition of the invention may of course include more than one compound of the invention.

In addition, the composition can comprise one or more additional active ingredients, for example compounds known to possess plant-growth regulant, herbicidal, fungicidal, insecticidal, acaricidal, antimicrobial or antibacterial properties. Alternatively the compound of the invention can be used in sequence with the other active ingredient.

The diluent or carrier in the composition of the invention can be a solid or a liquid optionally in association with a surface-active agent, for example a dispersing agent, emulsifying agent or wetting agent. Suitable surface-active agents include anionic compounds such as a carboxylate, for example a metal carboxylate of a long chain fatty acid; an *N*-acylsarcosinate; mono- or di-esters of phosphoric acid with fatty alcohol ethoxylates or alkyl phenol ethoxylates or salts of such esters; fatty alcohol sulfates such as sodium dodecyl sulfate, sodium octadecyl sulfate or sodium cetyl sulfate; ethoxylated fatty

alcohol sulfates; ethoxylated alkylphenol sulfates; lignin sulfonates; petroleum sulfonates; alkyl-aryl sulfonates such as alkyl-benzene sulfonates or lower alkylnaphthalene sulfonates, e.g. butyl-naphthalene sulfonate; salts of sulfonated naphthalene-formaldehyde condensates; salts of sulfonated phenol-formaldehyde condensates; or more complex sulfonates such as the amide sulfonates, e.g. the sulfonated condensation product of oleic acid and *N*-methyl taurine; the dialkyl sulfosuccinates, e.g. the sodium sulfonate of dioctyl succinate; acid derivatives of alkyl glycosides and alkylpolyglycosides materials and their metal salts, e.g. alkyl polyglycoside citrate or tartrate materials; or mono-, di- and tri-alkyl esters of citric acid and their metal salts.

10

Nonionic agents include condensation products of fatty acid esters, fatty alcohols, fatty acid amides or fatty-alkyl- or alkenyl-substituted phenols with ethylene and/or propylene oxide; fatty esters of polyhydric alcohol ethers, e.g. sorbitan fatty acid esters; condensation products of such esters with ethylene oxide, e.g. polyoxyethylene sorbitan fatty acid esters; alkyl glycosides, alkyl polyglycoside materials; block copolymers of ethylene oxide and propylene oxide; acetylenic glycols such as 2,4,7,9-tetramethyl-5-decyne-4,7-diol, ethoxylated acetylenic glycols; acrylic based graft copolymers; alkoxyated siloxane surfactants; or imidazoline type surfactants, e.g. 1-hydroxyethyl-2-alkylimidazoline.

15

Examples of a cationic surface-active agent include, for instance, an aliphatic mono-, di-, or polyamine as an acetate, naphthenate or oleate; an oxygen-containing amine such as an amine oxide, polyoxyethylene alkylamine or polyoxypropylene alkylamine; an amide-linked amine prepared by the condensation of a carboxylic acid with a di- or polyamine; or a quaternary ammonium salt.

20

The compositions of the invention can take any form known in the art for the formulation of agrochemicals, for example, a solution, an aerosol, a dispersion, an aqueous emulsion, a microemulsion, a dispersible concentrate, a dusting powder, a seed dressing, a fumigant, a smoke, a dispersible powder, an emulsifiable concentrate, granules or an impregnated strip. Moreover it can be in a suitable form for direct application or as a concentrate or primary composition which requires dilution with a suitable quantity of water or other diluent before application.

25

30

A dispersible concentrate comprises a compound of the invention dissolved in one or more water miscible or semi-water miscible solvents together with one or more surface active and/or polymeric material. Addition of the formulation to water results in the crystallisation of the active ingredient, the process being controlled by the surfactants and/or polymers resulting in a fine dispersion.

A dusting powder comprises a compound of the invention intimately mixed and ground with a solid pulverulent diluent, for example, kaolin.

An emulsifiable concentrate comprises a compound of the invention dissolved in a water-immiscible solvent which forms an emulsion or microemulsion on addition to water in the presence of an emulsifying agent.

A granular solid comprises a compound of the invention associated with similar diluents to those that may be employed in dusting powders, but the mixture is granulated by known methods. Alternatively it comprises the active ingredient absorbed or coated on a pre-formed granular carrier, for example, Fuller's earth, attapulgite, silica or limestone grit.

Wettable powders, granules or grains usually comprise the active ingredient in admixture with suitable surfactants and an inert powder diluent such as clay or diatomaceous earth.

Another suitable concentrate is a flowable suspension concentrate which is formed by grinding the compound with water or other liquid, surfactants and a suspending agent.

The concentration of the active ingredient in the composition of the present invention, as applied to plants is preferably within the range of 0.0001 to 1.0 per cent by weight, especially 0.0001 to 0.01 per cent by weight. In a primary composition, the amount of active ingredient can vary widely and can be, for example, from 5 to 95 per cent by weight of the composition.

The invention is generally applied to seeds, plants or their habitat. Thus, the compound can be applied directly to the soil before, at or after drilling so that the presence of active compound in the soil can control the growth of fungi which may attack seeds. When the soil is treated directly the active compound can be applied in any manner which allows it to



be intimately mixed with the soil such as by spraying, by broadcasting a solid form of granules, or by applying the active ingredient at the same time as drilling by inserting it in the same drill as the seeds. A suitable application rate is within the range of from 5 to 1000 g per hectare, more preferably from 10 to 500 g per hectare.

5

Alternatively the active compound can be applied directly to the plant by, for example, spraying or dusting either at the time when the fungus has begun to appear on the plant or before the appearance of fungus as a protective measure. In both such cases the preferred mode of application is by foliar spraying. It is generally important to obtain good control of  
10 fungi in the early stages of plant growth, as this is the time when the plant can be most severely damaged. The spray or dust can conveniently contain a pre- or post-emergence herbicide if this is thought necessary. Sometimes, it is practicable to treat the roots, bulbs, tubers or other vegetative propagule of a plant before or during planting, for example, by dipping the roots in a suitable liquid or solid composition. When the active compound is  
15 applied directly to the plant a suitable rate of application is from 0.025 to 5 kg per hectare, preferably from 0.05 to 1 kg per hectare.

In addition, the compounds of the invention can be applied to harvested fruits, vegetables or seeds to prevent infection during storage.

20


In addition, the compounds of the invention can be applied to plants or parts thereof which have been genetically modified to exhibit a trait such as fungal and/or herbicidal resistance.

In addition the compounds of the invention can be used to treat fungal infestations in timber  
25 and in public health applications. Also the compounds of the invention can be used to treat insect and fungus infestations in domestic and farm animals.

Compounds of the invention may be prepared, in known manner, in a variety of ways.


30

Compounds of formula Iai, i.e. compounds of general formula I where L is  $-\text{CH}(\text{R}^1)\text{NHCH}(\text{R}^2)-$ , may be prepared according to reaction scheme 1. Compounds of formula II or their hydrochloride salts can be condensed with compounds of formula III and the intermediate reduced with a suitable reagent such as sodium cyanoborohydride to give compounds of formula Iai.

Scheme 1


Compounds of formula II may be prepared by methods described in international application  
 5 PCT/GB/99/00304.

Compounds of formula Iai may also be prepared by reacting compounds of formula IV with  
 compounds of formula V in the same manner as above (Scheme 2).

Scheme 2

10


Compounds of formula Iaii, i.e. compounds of general formula I where L is  
 $-\text{CH(R}^1\text{)N(R}^3\text{)CH(R}^2\text{)-}$  and  $\text{R}^3$  is not hydrogen, may be prepared by reacting compounds of  
 formula Iai with a base and  $\text{R}^3\text{Q}$ , where Q is a leaving group such as a halogen. A suitable  
 base is triethylamine (Scheme 3).

15 Scheme 3

Compounds of formula Ib, i.e. compounds of general formula I where L is  
 $-\text{N(R}^3\text{)N(R}^4\text{)C(=X)-}$ , may be prepared according to reaction scheme 4 by reacting

compounds of formula VI with compounds of formula VII, where Q is a leaving group such as halogen, preferably chlorine. A preferred base is triethylamine.

Scheme 4




5

Compounds of formula Ic, i.e. compounds of general formula I where L is

-C(=O)N(R<sup>3</sup>)CH(R<sup>1</sup>)-, may be prepared by radical bromination of compounds of formula VIII, followed by reaction of these intermediates with compounds of formula IX according to scheme 5. Preferred reaction conditions are irradiation of a solution of VIII in carbon tetrachloride in the presence of *N*-bromosuccinimide and a catalytic amount of 2,2'-azobisisobutyronitrile, followed by addition of IX..

10

Scheme 5




15

Compounds of formula Id, i.e. compounds of general formula I where L is

-CH(R<sup>1</sup>)O(C=O)-, may be prepared according to reaction scheme 6 by formation of the cesium salt of compounds of formula XI, followed by reaction with compounds of formula X where Q is a suitable leaving group, such as chlorine.


Scheme 6



20


Compounds of formula Ie, i.e. compounds of general formula I where L is  
 $-\text{CH}(\text{R}^1)\text{OCH}(\text{R}^2)-$ , may be prepared by reaction of compounds of formula XII with a  
 suitable base such as sodium hydride, followed by reaction of the resulting anion with  
 compounds of formula X, where Q is a suitable leaving group such as halogen, according to  
 5 reaction scheme 7.

Scheme 7




Compounds of formula If, i.e. compounds of general formula I where L is  
 $-\text{N}(\text{R}^3)\text{C}(=\text{X})\text{N}(\text{R}^4)-$  and X is O or S, may be prepared according to reaction scheme 8 by  
 10 reaction of compounds of formula XIII with compounds of formula XIV, where X is O or S,  
 followed by the addition of compounds of formula XV. The order of addition of compounds  
 of formulae XIII and XV may be reversed.

Scheme 8




15

Compounds of formula Ig, i.e. compounds of general formula I where L is  
 $-\text{C}(\text{R}^1)=\text{C}(\text{R}^2)\text{C}(=\text{O})-$ , may be prepared according to reaction scheme 9 by reaction of  
 compounds of formula XVI with compounds of formula XVII in the presence of sodium  
 20 hydroxide.


Scheme 9

Compounds of formula Ih, i.e. compounds of general formula I where L is

- 5 - $\text{C}(\text{R}^1)=\text{N}-\text{N}(\text{R}^3)-$ , may be prepared by reacting compounds of formula XVIII with compounds of formula XIX according to reaction scheme 10.


Scheme 10

- 10 Compounds of formula Ii, i.e. compounds of general formula I where L is
- $\text{CH}(\text{R}^1)\text{N}=\text{C}(\text{R}^2)-$ , may be prepared according to reaction scheme 11 by reacting compounds of formula XX with a base, followed by reaction with compounds of formula XXI, where Q is a suitable leaving group, preferably chlorine. A suitable base is sodium hydride. Compounds of formula XX are known or can be prepared in a known manner by a
- 15 skilled chemist.

Scheme 11


Compounds of formula Ij, i.e. compounds of general formula I where L is  $-O-N=C(R^1)-$ , may be prepared according to reaction scheme 12 by the reaction of compounds of formula XXII with a base, followed by reaction with compounds of formula XXI, where Q is a suitable leaving group, preferably chlorine. A suitable base is sodium hydride.

5 Scheme 12




Compounds of formula XXII may be prepared according to reaction scheme 13.

Scheme 13




Compounds of formula Imi, i.e. compounds of general formula I where L is  $-O-NHC(=O)-$ , may be prepared according to reaction scheme 14 by the reaction of compounds of formula XXIII with compounds of formula XXIV, where Q is a suitable leaving group.

15 Scheme 14




Compounds of formula XXIV can be prepared from the corresponding hydroxy compounds by methods known to the skilled chemist. Compounds of formula XXIV can be isolated and used according to scheme 14 or generated *in situ* and used without isolation. A typical method, known to the skilled chemist, uses carbonyldiimidazole to generate compounds of formula XXIV *in situ*.

Compounds of formula XXIII can be prepared according to reaction scheme 15.

Scheme 15

Compounds of formula Imii, i.e. compounds of general formula I where L is  
 5 -O-N(R<sup>3</sup>)C(=O)- wherein R<sup>3</sup> is not hydrogen, may be prepared by reaction of compounds of formula Imi with a base, followed by reaction with R<sup>3</sup>Q, where Q is a suitable leaving group, such as a halogen. A suitable base is potassium *tert*-butoxide (Scheme 16).

Scheme 16

10

Other methods will be apparent to the chemist skilled in the art, as will be the methods for preparing starting materials and intermediates.

15 Collections of compounds of formula I may also be prepared in a parallel manner, either manually, automatically or semi-automatically. This parallel preparation may be applied to the reaction procedure, work-up or purification of products or intermediates. For a review of such procedures see by S.H. DeWitt in "Annual Reports in Combinatorial Chemistry and Molecular Diversity: Automated synthesis", Volume 1, Verlag Escom 1997, pages 69 to 77.

20 Furthermore, compounds of the formula I may be prepared using solid-supported methods, where the reactants are bound to a synthetic resin. See for example: Barry A. Bunin in "The

Combinatorial Index", Academic Press, 1998 and "The tea-bag method" (Houghten, US 4,631,211; Houghten et al., Proc. Natl. Acad. Sci, 1985, 82, 5131-5135).

The invention is illustrated in the following Examples. Structures of isolated, novel  
5 compounds were confirmed by NMR and/or other appropriate analyses.

#### Example 1

*N*-(2-Chlorobenzyl)-*N*-{1-[3-chloro-5-(trifluoromethyl)-2-pyridyl]ethyl}amine  
(Compound 27)

10  $\alpha$ -Methyl-[3-chloro-5-(trifluoromethyl)-2-pyridyl]methylamine (0.2 g) was dissolved in trimethylorthoformate (10 ml) and triethylamine (0.22 ml) was added. After 5 minutes 2-chlorobenzaldehyde (0.26 g) was added and the resulting mixture stirred for 3.5 hours at room temperature to give a precipitate. Sodium cyanoborohydride (1.5 ml, 0.1M solution in tetrahydrofuran) and acetic acid (0.1 ml) were then added and the mixture stirred for 16  
15 hours at room temperature. Brine (5 ml) and water (10 ml) were then added and the mixture stirred for 20 minutes. The phases were separated and the organic phase evaporated. The residue was purified by silica gel chromatography to give the title product, m.p. 117 °C.

#### Example 2

20 *N*-{[3-Chloro-5-(trifluoromethyl)-2-pyridyl]methyl}-*N*-[1-(3,4-difluorophenyl)-ethyl]amine  
(Compound 23)


Triethylamine (0.08 ml) and 1-(3,4-difluorophenyl)-1-ethanamine (0.11 g) were dissolved in trimethylorthoformate (10 ml). 3-Chloro-(5-trifluoromethyl)-pyridine-2-carboxaldehyde (0.15 g) was added and the solution stirred for 4 hours at room temperature. Sodium  
25 cyanoborohydride (1 ml, 0.1M solution in tetrahydrofuran) and acetic acid (0.07 ml) were then added and the mixture stirred for 16 hours at room temperature. Brine (5 ml) and water (10 ml) were then added and the mixture stirred for 20 minutes. The phases were separated and the organic phase evaporated. The crude material was purified by silica gel  
chromatography to give the title product, <sup>1</sup>H N.M.R (CDCl<sub>3</sub>)  $\delta$ (ppm) 1.4 (3H, d), 2.6 (1H,  
30 broad s), 3.8 (1H, q), 3.9 (1H, s), 7.1–7.3 (3H, m), 7.9 (1H, s) and 8.8 (1H, s).



Example 3Ethyl 2-[acetyl(benzyl)amino]-2-[3-chloro-5-(trifluoromethyl)-2-pyridyl]acetate  
(Compound 4)

To a solution of compound 1 (0.6 mmol) in diethyl ether was added triethylamine (0.7 mmol) followed by acetyl chloride in diethyl ether (0.7 mmol). The mixture was stirred for two hours at room temperature before the addition of hydrochloric acid (8 ml, 2M). The organic phase was isolated, washed with sodium bicarbonate (10ml), dried over magnesium sulfate and evaporated to yield the title compound, <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) (ppm) δ1.2 (3H, t), 2.3 (3H, s), 4.2 (2H, q), 4.8 (2H, q), 6.8-7.1 (6H, m), 7.5 (1H, s) and 8.5 (1H, s).



The following compounds of formula Iz (see Table A), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -CH(R<sup>1</sup>)N(R<sup>3</sup>)CH(R<sup>2</sup>)-, may be prepared by methods analogous to those of Examples 1, 2 and 3. The amine starting materials were obtained using methods described in international application PCT/GB/99/00304.




(Iz)

Table A

Cmp	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	A <sup>2</sup>	m.p.°C
1	EtOC(=O)-	H	H	phenyl	oil
2	EtOC(=O)-	H	H	2-Cl-phenyl	oil
3	EtOC(=O)-	H	H	3,4-methylenedioxyphenyl	oil
4	EtOC(=O)-	H	MeC(=O)-	phenyl	oil
5	EtOC(=O)-	H	MeOCH <sub>2</sub> C(=O)-	phenyl	oil
6	EtOC(=O)-	H	MeOC(=O)-C(=O)-	phenyl	104-7
7	EtOC(=O)-	H	MeC(=O)-	2-Cl-phenyl	77-81
8	EtOC(=O)-	H	MeOCH <sub>2</sub> C(=O)-	2-Cl-phenyl	oil
9	EtOC(=O)-	H	MeOC(=O)-C(=O)-	2-Cl-phenyl	128-31
10	EtOC(=O)-	H	MeC(=O)-	3,4-methylenedioxyphenyl	oil

Cmp	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	A <sup>2</sup>	m.p./°C
11	EtOC(=O)-	H	MeOCH <sub>2</sub> C(=O)-	3,4-methylenedioxyphenyl	oil
12	EtOC(=O)-	H	MeOC(=O)-C(=O)-	3,4-methylenedioxyphenyl	106-9
13	H	MeOC(=O)CH <sub>2</sub> -	H	phenyl	oil
14	H	EtOC(=O)CH <sub>2</sub> -	H	phenyl	oil
15	H	H	H	phenyl	oil
16	H	H	H	2-Cl-6-F-phenyl	oil
17	H	Me	H	2-Cl-phenyl	oil
18	H	Me	H	2,6-diF-phenyl	oil
19	H	Me	H		oil
20	H	Me	H	4-tolyl	oil
21	H	H	H	2,5-diF-phenyl	oil
22	H	Me	H	4-NO <sub>2</sub> -phenyl	oil
23	H	Me	H	3,4-diF-phenyl	oil
24	H	H	H	2-Cl-phenyl	oil
25	H	H	H	4-PhO-phenyl	oil
26	H	H	H	2-NO <sub>2</sub> -phenyl	oil
27	Me	H	H	2-Cl-phenyl	117
28	Me	H	H	2-NO <sub>2</sub> -phenyl	136
29	H	Me	H	phenyl	oil
30	H	Me	H	3-CF <sub>3</sub> O-phenyl	oil
31	H	Me	H	4-CF <sub>3</sub> O-phenyl	oil
32	H	Me	H		oil
33	H	Me	H	4-Cl-phenyl	oil
34	H	Me	H	4-Br-phenyl	oil
35	Me	H	H	cyclohexyl	oil
36	Me	H	H	2-F-phenyl	oil
37	Me	H	H	4-Cl-phenyl	oil
38	Me	H	H	2,5-diMeO-phenyl	oil

Cmp	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	A <sup>2</sup>	m.p.°C
39	Me	H	H	2-Cl-6-F-phenyl	oil
40	Me	H	H	2-Br-phenyl	oil
41	Me	H	H	3-CF <sub>3</sub> O-phenyl	oil
42	Me	H	H	4-MeS-phenyl	oil
43	Me	H	H	2,5-xylyl	oil
44	H	H	H	cyclohexyl	oil
45	H	H	H	3-Br-phenyl	oil
46	H	H	H	4-Me <sub>2</sub> N-phenyl	oil
47	H	H	H	4-Cl-phenyl	oil
48	H	H	H	2-F-phenyl	oil
49	H	H	H	2,5-diMeO-phenyl	oil
50	H	H	H	2-Br-phenyl	oil
51	H	H	H	4-NO <sub>2</sub> -phenyl	oil
52	H	H	H	2,5-xylyl	oil
53	H	Me	H		oil
54	H	H	H	pentaF-phenyl	oil

The <sup>1</sup>H N.M.R. or mass spectral data of those compounds in Table A which were not solid at room temperature are presented below.

5 Compound 1

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 1.2 (3H, t), 2.9 (1H, broad s), 3.8 (1H, q), 4.2 (2H, m), 5.0 (1H, s), 7.4-7.2 (5H, m), 7.9 (1H, s), 8.7 (1H, s).

Compound 2

10 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 1.2 (3H, t), 3.1 (1H, broad s), 4.0 (2H, q), 4.2 (2H, m), 5.1 (1H, s), 7.5-7.2 (4H, m), 8.0 (1H, s), 8.7 (1H, s).

Compound 3

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 1.2 (3H, t), 2.4 (1H, broad s), 3.8 (2H, q), 4.2 (2H, m), 5.0 (1H, s), 5.9 (2H, s), 6.74 (1H, d), 6.76 (1H, s), 6.83 (1H, s), 7.9 (1H, s), 8.7 (1H, s).

5 Compound 4

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 1.2 (3H, t), 2.3 (3H, s), 4.2 (2H, q), 4.8 (2H, q), 6.8-7.1 (6H, m), 7.5 (1H, s), 8.5 (1H, s).

Compound 5

10 *m/z* (APCI) 445 (M+H)<sup>+</sup>.

Compound 8

*m/z* (APCI) 479 (M+H)<sup>+</sup>.

15 Compound 10

*m/z* (APCI) 487 (M<sup>-</sup>)

Compound 11

*m/z* (APCI) 459 (M+H)<sup>+</sup>.

20

Compound 13

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 2.7 (1H, dd), 2.9 (1H, dd), 3.6 (3H, s), 3.9 (2H, s), 4.2 (1H, m), 7.3 (5H, m), 7.8 (1H, s), 8.8 (1H, m).

25 Compound 14

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.2 (3H, t), 2.7 (1H, dd), 2.8 (1H, dd), 3.9 (2H, s), 4.1 (2H, q), 4.2 (1H, m), 7.2-7.4 (5H, m), 7.8 (1H, s), 8.8 (1H, s).

Compound 15

30 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 4.2 (2H, s), 5.3 (2H, s), 7.3 (6H, m), 8.8 (1H, s), 8.9 (1H, s).

Compound 16

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 2.7 (1H, broad s), 4.05 (4H, s), 6.9-7.2 (3H, m), 7.8 (1H, s), 8.6 (1H, s).

5 Compound 17

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.5 (1H, broad s), 3.9 (2H, m), 4.3 (1H, q), 7.1-7.3 (3H, m) 7.5 (1H, m) 7.8 (1H, s), 8.7 (1H, s).

Compound 18

10 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.5 (3H, d), 2.6 (1H, broad s), 3.8 (1H, m), 4.0 (1H, m) 4.3 (1H, q), 6.8 (2H, m) 7.1 (1H, m) 7.8 (1H, s) 8.6 (1H, s).

Compound 19

15 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d) 2.6 (1H, broad s), 3.8 (1H, q), 4.0 (2H, s), 4.3 (4H, s) 6.8 (2H, s), 6.9 (1H, s), 7.9 (1H, s), 8.7 (1H, s).

Compound 20

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.5 (3H, d), 2.4 (3H, s), 2.8 (1H, broad s), 3.8 (1H, q), 4.0 (2H, m), 7.2 (2H, d), 7.3 (2H, d), 7.8 (1H, s), 8.7 (1H, s).

20

Compound 21

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 2.5 (1H, broad s), 4.0 (2H, s), 4.1 (2H, s), 6.8 (2H, m), 7.2 (1H, m), 7.8 (1H, s), 8.6 (1H, s).

25 Compound 22

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.5 (1H, broad s), 3.86 (2H, s), 3.9 (1H, m), 7.5 (2H, d), 7.8 (1H, s), 8.1 (2H, d), 8.7 (1H, s).

Compound 23

30 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.6 (1H, broad s), 3.8 (1H, q), 3.9 (1H, s), 7.1-7.3 (3H, m), 7.9 (1H, s), 8.8 (1H, s).

Compound 24

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 4.1 (4H, m), 4.4 (1H, dd), 4.6 (1H, dd), 6.5 (1H, broad s), 7.2-7.4 (5H, m), 7.8 (1H, s), 8.6 (1H, s).

5

Compound 25

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 3.9 (1H, dd), 4.2 (1H, dd), 4.4 (2H, m), 6.0 (1H, broad s), 6.9-7.0 (5H, m), 7.2-7.4 (4H, m), 8.0 (1H, s), 8.7 (1H, s).

10. Compound 26

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 4.3 (2H, m), 4.7 (2H, m), 7.0 (1H, broad s), 7.6 (1H, m), 7.7 (2H, m), 7.9 (1H, s), 8.2 (1H, m), 8.6 (1H, s).

Compound 29

15 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.5 (3H, s), 2.6 (1H, broad s), 3.9 (1H, q), 4.0 (2H, s), 7.2-7.4 (5H, m), 7.8 (1H, s), 8.7 (1H, s).

Compound 30

20 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.6 (1H, broad s), 3.8 (1H, q), 3.9 (2H, s), 4.0 (2H, s), 7.0 (1H, m), 7.2-7.3 (3H, m), 7.8 (1H, s), 8.7 (1H, s).

Compound 31

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.5 (3H, d), 2.7 (1H, broad s), 3.8 (1H, q), 6.5 (1H, m), 7.1 (2H, d), 7.4 (2H, d), 7.9 (1H, s), 8.8 (1H, s).

25

Compound 32

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.5 (3H, d), 1.8 (4H, m), 2.8 (4H, m), 3.8 (1H, q), 4.0 (2H, s), 7.1 (3H, m), 7.9 (1H, s), 8.8 (1H, s).

Compound 33

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.42 (3H, d), 2.62 (1H, broad, s), 3.83 (1H, q), 3.92 (2H, s), 7.3 (4H, s), 7.82 (1H, s); 8.75 (1H, s).

5 Compound 34

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.42 (3H, d), 2.58 (1H, s, broad), 3.82 (1H, q), 3.92 (2H, s), 7.25 (2H, m), 7.45 (2H, m), 7.85 (1H, s), 8.72 (1H, s).

Compound 35

10 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 1.38 (3H, d), 4.35 (1H, q), 7.85 (1H, s), 8.75 (1H, s).

Compound 36

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.38 (3H, d), 2.35 (1H, broad, s), 3.68 (2H, m), 4.42 (1H, q),  
15 6.95 (1H, m), 7.05 (1H, m), 7.16 (1H, m), 7.32 (1H, m), 7.82 (1H, s), 8.75 (1H, s).

Compound 37

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 1.38 (3H, d), 3.56 (2H, m), 4.4 (1H, q), 7.88 (1H, s), 8.78 (1H, s).

20

Compound 39

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.3 (3H, d), 2.7 (1H, broad, s), 3.8 (2H, s), 4.4 (1H, q),  
6.75 (1H, m), 6.98 (2H, m), 7.72 (1H, s), 8.55 (1H, s).

25 Compound 40

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.41 (3H, d), 2.3 (1H, broad, s), 3.72 (2H, m), 4.25 (1H, q),  
7.05-7.5 (4H, m), 7.85 (1H, s), 8.75 (1H, s).

Compound 41

30 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.38 (3H, d), 2.4 (1H, s, broad), 3.6 (2H, m), 4.4 (1H, q),  
7.05-7.5 (4H, m), 7.88 (1H, s), 8.78 (1H, s).

Compound 42

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.38 (3H, d), 2.48 (3H, s), 3.6 (2H, m), 4.45 (1H, q),  
7.2 (4H, m), 7.88 (1H, s), 8.78 (1H, s).

5

Compound 43

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.25 (3H, s), 2.32 (3H, s), 2.5 (1H, broad, s),  
3.58 (2H, m), 4.48 (1H, q), 6.9-7.08 (3H, m), 7.9 (1H, s), 8.78 (1H, s).

10 Compound 44

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 2.53 (2H, d), 4.1 (2H, s), 7.85 (1H, s),  
8.75 (1H, s).

Compound 45

15 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 3.85 (1H, s), 4.1 (1H, s), 7.9 (1H, s), 8.75 (1H, s).

Compound 46

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 7.8 (1H, s), 8.68 (1H, s), 3.5 (1H, m), 3.9 (2H,  
m), 4.1 (1H, m).

20

Compound 47

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 3.86 (2H, s), 4.08 (2H, s), 7.90 (1H, s),  
8.72 (1H, s).

25 Compound 48

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 2.65 (1H, s, broad), 3.98 (2H, s), 4.15 (2H, s),  
7.9 (1H, s), 8.75 (1H, s).

Compound 49

30 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 3.80 (3H, s), 3.85 (3H, s), 6.88 (2H, m),  
7.06 (1H, m), 7.90 (1H, s), 8.72 (1H, s).



Compound 50

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 4.0 (2H, s), 4.1 (2H, s), 7.85 (1H, s), 8.70 (1H, s).

5 Compound 51

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 2.65 (1H, broad, s), 3.83 (2H, s), 4.0 (2H, s), 7.8 (1H, s), 8.65 (1H, s).

Compound 52

10 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 3.83 (2H, s), 4.18 (2H, s), 7.9 (1H, s), 8.75 (1H, s).

Compound 53

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 1.4 (3H, d), 2.5 (1H, broad s), 3.8 (1H, q), 3.9 (2H, s), 6.0 (2H, 15 s), 6.8-6.9 (3H, m), 7.9 (1H, s), 8.7 (1H, s).

Compound 54


<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) selected peaks at 7.75 (1H, s), 8.80 (1H, s).

20 Example 4

N'-1-[3-Chloro-5-(trifluoromethyl)-2-pyridyl]-2,6-dichloro-1-benzenecarbohydrazide  
(Compound 102)

3-Chloro-5-(trifluoromethyl)pyrid-2-ylhydrazine (0.32 g) was dissolved in dichloromethane (7 ml) and treated dropwise with 2,6-dichlorobenzoyl chloride (0.31 g) in dichloromethane 25 (2 ml). Triethylamine (0.15 g) was then added and the reaction stirred at room temperature overnight. The organic solution was washed sequentially with sodium bicarbonate solution and brine and evaporated to give a solid. The residue was purified by trituration (dichloromethane) to give the title compound, m.p. 212-5 °C.

30 The following compounds of formula Iy (see Table B), i.e. compounds of general formula I where L is -N(R<sup>3</sup>)NHC(=O)-, may be prepared by methods analogous to those of Example 4.



(ly)


Table B

Cmp	A <sup>1</sup>	R <sup>3</sup>	A <sup>2</sup>	m.p./°C
101	3-Cl-5-CF <sub>3</sub> -phenyl	H	2-Cl-phenyl	168-70
102	3-Cl-5-CF <sub>3</sub> -phenyl	H	2,6-diCl-phenyl	212-5
103	3-Cl-5-CF <sub>3</sub> -phenyl	H	2-NO <sub>2</sub> -phenyl	182-3
104	3-Cl-5-CF <sub>3</sub> -phenyl	H	2,6-diMeO-phenyl	204-6
105	3-Cl-5-CF <sub>3</sub> -phenyl	H	2-tolyl	168-9
106	3-Cl-5-CF <sub>3</sub> -phenyl	H	4,6-diMeO-pyrimidin-2-yl	170-1
107	3-Cl-5-CF <sub>3</sub> -phenyl	H	cyclopropyl	152-4
108	3-Cl-5-CF <sub>3</sub> -phenyl	H	cyclohexyl	111-4
109	3-Cl-5-CF <sub>3</sub> -phenyl	Me	2,6-diCl-phenyl	219-20
110	3-Cl-5-CF <sub>3</sub> -phenyl	Me	2-NO <sub>2</sub> -phenyl	198-9
111	3-Cl-5-CF <sub>3</sub> -phenyl	Me	2,6-diMeO-phenyl	234-6
112	3-Cl-5-CF <sub>3</sub> -phenyl	Me	2-tolyl	202-4
113	3-Cl-5-CF <sub>3</sub> -phenyl	Me	2-Cl-6-F-phenyl	207-8
114	3-Cl-5-CF <sub>3</sub> -phenyl	Me	4,6-diMeO-pyrimidin-2-yl	178-80
115	3-Cl-5-CF <sub>3</sub> -phenyl	Me	cyclopropyl	159-60
116	3-Cl-5-CF <sub>3</sub> -phenyl	Me	cyclohexyl	216-9
117	5-Cl-3-CF <sub>3</sub> -phenyl	H	2,6-diCl-phenyl	199-203
118	5-Cl-3-CF <sub>3</sub> -phenyl	H	2-NO <sub>2</sub> -phenyl	156-8
119	5-Cl-3-CF <sub>3</sub> -phenyl	H	2,6-diMeO-phenyl	194-5
120	5-Cl-3-CF <sub>3</sub> -phenyl	H	2-tolyl	180-1
121	5-Cl-3-CF <sub>3</sub> -phenyl	H	2-Cl-6-F-phenyl	173-5
122	5-Cl-3-CF <sub>3</sub> -phenyl	H	4,6-diMeO-pyrimidin-2-yl	158
123	5-Cl-3-CF <sub>3</sub> -phenyl	H	cyclopropyl	143-5
124	5-Cl-3-CF <sub>3</sub> -phenyl	H	cyclohexyl	121

Cmp	A <sup>1</sup>	R <sup>3</sup>	A <sup>2</sup>	m.p./°C
125	3-Cl-5-CF <sub>3</sub> -phenyl	H	2,3,6-triF-phenyl	154-6
126	3-Cl-5-CF <sub>3</sub> -phenyl	H	2-Cl-6-F-phenyl	192

Example 5*N*-2-(Phenylethyl)-3-chloro-5-(trifluoromethyl)-2-pyridinecarboxamide(Compound 206)




- 5 3-Chloro-(5-trifluoromethyl)pyridine-2-carboxaldehyde (0.15 g) was dissolved in carbon tetrachloride (10 ml). 2,2'-Azobisisobutyronitrile (0.002 g) and *N*-bromosuccinimide (0.16 g) were added and the mixture was heated to reflux using a sun lamp. After 45 minutes the solution was cooled down to 0 °C. (R)-(+)- $\alpha$ -methylbenzylamine (0.09 g) in carbon tetrachloride (0.3 ml) was added and stirred for 20 minutes at 0°C, then for 3 hours at room
- 10 temperature. The mixture was diluted with dichloromethane and washed with water. The organic layer was isolated, dried over magnesium sulphate and evaporated to give the crude compound. The crude material was purified by silica gel chromatography to give the title product, m.p. 88 °C.
- 15 The following compounds of formula Ix (see Table C), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -C(=O)NHCH(R<sup>1</sup>)-, may be prepared by methods analogous to those of Example 5.



(Ix)

Table C


Cmp	R <sup>1</sup>	A <sup>2</sup>	m.p.(°C)
201	H	2,6-diF-phenyl	137
202	Me	2,6-diF-phenyl	97
203	H	2-Cl-phenyl	100-7
204	H	2,6-diCl-phenyl	114-6

Cmp	R <sup>1</sup>	A <sup>2</sup>	m.p.(°C)
205	Me	2-Cl-phenyl	120
206	Me	phenyl	88
207	Me	4-Cl-phenyl	129
208	Me	4-Br-phenyl	139
209	Me	3,4-diF-phenyl	127
210	Me		123
211	Me	4-CF <sub>3</sub> O-phenyl	95
212	Me	3-CF <sub>3</sub> O-phenyl	114
213	Me		125
214	Me		129
215	H	4-tolyl	113

**Example 6**[3-Chloro-5-(trifluoromethyl)-2-pyridyl]methyl 2-chlorobenzoateCompound 301

- 5 To a solution of 2-chlorobenzoic acid (0.1 g) in dimethylformamide was added cesium carbonate (0.1 g) and the resulting solution was stirred for 1 hour. 3-Chloro-2-(chloromethyl)-5-trifluoromethyl pyridine (0.14 g) was added and stirring was continued for a further 48 hours. The solution was diluted with diethyl ether (10 ml) and washed with water (10 ml). The organic phase was separated, dried and evaporated to give a crude
- 10 product. Silica gel chromatography (petrol/diethyl ether 7:3) gave the title compound, <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 5.6 (2H, s), 7.3 (1H, m), 7.4 (2H, m) 7.87 (1H, s), 7.88 (1H, d) and 8.8 (1H, s).

The following compounds of formula Iw (see Table D), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -CH<sub>2</sub>O(C=O)-, may be prepared by methods analogous to those of Example 6.



(Iw)

Table D

Cmp	A <sup>2</sup>	m.p./°C
301	2-Cl-phenyl	oil
302	2,6-diCl-phenyl	93-5


**Example 7**

[3-Chloro-5-(trifluoromethyl)-2-pyridyl]methyl(2,4-dichlorobenzyl) ether

**(Compound 401)**

To a solution of 2,4-dichlorobenzyl alcohol (0.27 g) in tetrahydrofuran under nitrogen was added sodium hydride (1.1 equivalents) portionwise. The resulting solution was stirred at room temperature for 1 hour before the addition of 3-chloro-2-(chloromethyl)-5-trifluoromethyl pyridine (0.35 g) in tetrahydrofuran dropwise. The solution was then stirred at room temperature for 16 hours. The solution was treated with a tetrahydrofuran/methanol solution and the solvent then evaporated. The residue was partitioned between water and ethyl acetate, the organic phase was isolated, washed with brine, dried and evaporated to yield the crude product. Silica gel chromatography (petrol/ethyl acetate 95:5) furnished the title compound, <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ(ppm) 4.8 (2H, s), 4.9 (2H, s), 7.3 (1H, m), 7.4 (1H, m), 7.5 (1H, m) 8.0 (1H, s) and 8.8 (1H, m).

The following compounds of formula Iv (see Table E), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -CH<sub>2</sub>OCH<sub>2</sub>-, may be prepared by methods analogous to those of Example 7.



(iv)

Table E

Cmp	A <sup>2</sup>	m.p./°C
401	2,4-diCl-phenyl	oil
402	2,6-diCl-phenyl	oil

The <sup>1</sup>H N.M.R. data of those compounds in Table E which were not solid at room temperature are presented below.

Compound 402


<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 4.9 (2H, s), 5.0 (2H, s), 7.2 (1H, m), 7.3 (2H, m), 8.0 (1H, s), 8.8 (1H, s).

Example 8

N-[2-Chloro-5-(trifluoromethyl)-2-pyridyl]-N'-(2,6-dichlorophenyl)urea  
(Compound 501)

A solution of triphosgene (1.1 g) in dichloromethane (20 ml) was added over 30 minutes at room temperature to a stirred solution of 2-amino-3-chloro-5-(trifluoromethyl)pyridine (1.96 g) and triethylamine (2 ml) in dichloromethane (35 ml). After 15 minutes a solution of 2,6-dichloroaniline (1.62 g) and triethylamine (2 ml) in dichloromethane (20 ml) was added rapidly and the resulting mixture stirred for 30 minutes before solvent evaporation. The residue was suspended in ethyl acetate and the solid filtered off. The filtrate was washed with potassium hydrogen sulfate solution, sodium bicarbonate solution and then brine. Drying (MgSO<sub>4</sub>) and solvent evaporation yielded the crude product, which was purified by silica gel chromatography to give the title compound, m.p. 155-8 °C.

The following compounds of formula Iu (see Table F), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -NHC(=O)NH-, may be prepared by methods analogous to those of Example 8.



(Iu)

Table F

Cmp	A <sup>2</sup>	m.p./°C
501	2,6-diCl-phenyl	155-8
502	phenyl	173-5
503	2-NO <sub>2</sub> -phenyl	178-80

#### 5 Example 9

##### 3-[3-Chloro-5-(trifluoromethyl)-2-pyridyl]-1-(2-nitrophenyl)-2-propen-1-one (Compound 601)

Sodium hydroxide (0.55 g) was dissolved in water (5 ml) and the resulting solution was diluted with ethanol (3 ml). 2-Nitroacetophenone (1.8 g) was added at 20°C, and the solution was stirred for 5 minutes. 3-Chloro-5-(trifluoromethyl)pyridine-2-carboxaldehyde (2.25 g) was added and stirring was continued for 16 hours. The solution was acidified with acetic acid, the organic layer separated, dried over magnesium sulfate, filtered and evaporated to give a brown oil. Silica gel column chromatography, followed by recrystallisation (petrol) afforded the title compound, 88-9 °C.

15

#### Example 10

##### 3-Chloro-5-(trifluoromethyl)-2-pyridinecarbaldehyde 2-(2-nitrophenyl)hydrazone (Compound 701)


A mixture of 3-chloro-5-(trifluoromethyl)pyridine-2-carboxaldehyde (1.05 g) and 2-nitrophenylhydrazine (0.76 g) in ethanol (75 ml) was heated at reflux for 2.5 hours and then allowed to cool to room temperature overnight. The resulting orange solid was isolated by filtration and recrystallised (petrol) to afford the title compound as a mixture of isomers, m.p. 127-35 °C.

20

Example 11[3-Chloro-5-(trifluoromethyl)-2-pyridyl][(diphenylmethylene)amino]methyl cyanide(Compound 803)

To a suspension of 60% sodium hydride (4.0 g) in dimethylformamide under a nitrogen atmosphere at 0°C was added a solution of [(diphenylmethylene)amino]methyl cyanide (11.1g) in dimethylformamide dropwise, whilst maintaining the temperature between 0°C and 2°C. The solution was stirred at 0°C for 1 hour. 2,3-Dichloro-5-trifluoromethylpyridine (7 ml) in dimethylformamide was added dropwise and the mixture stirred for 30 minutes at 0°C before warming to ambient temperature over 3 hours. The mixture was cooled to 10°C, ethanol (3 ml) added and the solution stirred for 15 minutes. The reaction mixture was then poured as a thin stream into a vigorously stirred mixture of diethyl ether (500ml) and ammonium chloride solution (500 ml). The organic layer was separated and washed with ammonium chloride solution (2x150 ml), dried, filtered and evaporated to give a residue. Silica gel chromatography (diethyl ether:petrol 5:95) gave the title product as a pale brown solid, m.p. 108-10 °C.

The following compounds of formula It (see Table G), i.e. compounds of general formula I where A<sup>1</sup> is 3-Cl-5-CF<sub>3</sub>-2-pyridyl, L is -CH(R<sup>1</sup>)N=C(Ph)-, and A<sup>2</sup> is phenyl may be prepared by methods analogous to those of Example 11.



(It)

Table G

Cmp	R <sup>1</sup>	m.p./°C
801	CH <sub>2</sub> CN	82-4
802	CO <sub>2</sub> Et	oil
803	CN	108-10



The mass spectral data of the compound in Table G which was not solid at room temperature is presented below.

Compound 802

5  $m/z$  (EI) 373 ( $M^+ - CO_2Et$ )

Example 12

1-Biphenyl-1-ethanone O-1-[3-chloro-5-(trifluoromethyl)-2-pyridyl] oxime

(Compound 936)

10 To 4-acetylbiphenyl oxime (2.5 g) in dimethylformamide (13 ml) under a nitrogen atmosphere was added sodium hydride (0.5 g) portionwise with cooling. The resulting mixture was stirred at 40°C for 20 minutes until the formation of a suspension occurred. 2,3-Dichloro-5-(trifluoromethyl)pyridine (2.5 g) in dimethylformamide (7 ml) was then added and the resulting mixture stirred for 18 hours at room temperature. The mixture was  
15 treated with isopropanol (2 ml) and stirred for 5 minutes before pouring into an ice water/brine solution (300 ml). The resulting precipitate was extracted with diethyl ether (2x125 ml), the organics washed with water, dried, filtered and evaporated to give a solid which on trituration (diethyl ether) and recrystallisation (toluene) yielded the title compound, m.p. 122 °C.

20

Preparation of Starting Material


4-Acetylbiphenyl Oxime

To a suspension of 4-acetylbiphenyl (25.4 g) in ethanol (230 ml) and water (4 ml) under a nitrogen atmosphere was added hydroxylamine hydrochloride (14.5 g) in  
25 water (25 ml) followed by 50% aqueous potassium hydroxide solution (40 g). The resulting mixture was heated at reflux for 18 hours and then cooled to room temperature. The mixture was added to ice/water (500 ml) and acidified to pH 2 to give a precipitate. The solid was filtered off, washed with water until the washings were at pH 6 and then recrystallised from ethanol to give the title compound.

30


The following compounds of formula Is (see Table H), i.e. compounds of general formula I where L is  $-O-N=C(R^1)-$ , may be prepared by methods analogous to those of Example 12.


The crossed bond in Is indicates that the compounds may exist as cis or trans isomers about the double bond. Isolation of both isomers was possible for some compounds.



(Is)

Table H

Cmp	A <sup>1</sup>	R <sup>1</sup>	A <sup>2</sup>	m.p.(°C)
901	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-Cl-phenyl	96-7
902	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	4-pyridyl	205-6
903	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	3-(2-Cl-4-CF <sub>3</sub> -phenoxy)phenyl	65-7
904	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2-Cl-6-F-phenyl	119-23
905	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2,6-diCl-phenyl	136-7
906	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	1-Me-2-pyrolyl	88-9
907	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-tolyl	oil
908	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-tolyl	oil
909	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	3-CF <sub>3</sub> -phenyl	oil
910	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-CF <sub>3</sub> -phenyl	oil
911	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me		oil
912	3-Cl-5-CF <sub>3</sub> -2-pyridyl	tBu	2-pyridyl	oil
913	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-thienyl	oil
914	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	4-MeO-phenyl	oil
915	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2,4-xylyl	oil
916	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	6-Me-2-pyridyl	oil
917	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-naphthyl	oil
918	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	1-naphthyl	oil
919	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	4-EtO-phenyl	oil
920	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2-tolyl	oil
921	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2-MeO-phenyl	oil

Cmp	A <sup>1</sup>	R <sup>1</sup>	A <sup>2</sup>	m.p.(°C)
922	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Et	phenyl	oil
923	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	3-NO <sub>2</sub> -phenyl	116-8
924	3-Cl-5-CF <sub>3</sub> -2-pyridyl	CF <sub>3</sub>	2-tolyl	oil
925	3-Cl-5-CF <sub>3</sub> -2-pyridyl	(EtO) <sub>2</sub> P(=O)-	cyclohexyl	oil
926	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	phenyl	76
927	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	phenyl	oil
928	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2-NO <sub>2</sub> -phenyl	oil
929	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	2-Cl-phenyl	87
930	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	3-tolyl	oil
931	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	3-pyridyl	oil
932	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	3-pyridyl	137-8
933	3-Cl-5-CF <sub>3</sub> -2-pyridyl	H	1-naphthyl	85-90
934	3,5-diCl-2-pyridyl	Me	2-Cl-phenyl	127
935	3,5-diCl-2-pyridyl	Me	2-Cl-phenyl	70-1
936	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	biphenyl	122
937	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2,6-diCl-phenyl	128-9
938	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2,6-diCl-phenyl	71-2
939	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2-CN-phenyl	139-43
940	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2-Cl-phenyl	83-4
941	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2-Cl-phenyl	88
942	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2-MeSO <sub>2</sub> -phenyl	oil
943	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Ph	2-naphthyl	oil
944	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	6-MeO-2-naphthyl	oil
945	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	4-F-1-naphthyl	oil
946	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	4-cyclohexyl-phenyl	oil
947	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me		oil
948	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Pr	4-Cl-phenyl	oil
949	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	cyclohexyl	oil

Cmp	A <sup>1</sup>	R <sup>1</sup>	A <sup>2</sup>	m.p.(°C)
950	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	4-PhO-phenyl	oil
951	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2,5-diMe-3-furyl	oil
952	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	3,5-diMe-isothiazol-4-yl	oil
953	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Et	2,4-diCl-phenyl	oil
954	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	2,3-diCl-phenyl	oil
955	3-Cl-5-CF <sub>3</sub> -2-pyridyl	-CN	2-pyridyl	oil
956	3-Cl-5-CF <sub>3</sub> -2-pyridyl	CF <sub>3</sub>	2-thienyl	oil
957	3-Cl-5-CF <sub>3</sub> -2-pyridyl	Me	4-pyridyl	oil
958	3-Cl-5-CF <sub>3</sub> -2-pyridyl	4-Cl-phenyl	4-Cl-phenyl	oil

The <sup>1</sup>H N.M.R or mass spectral data of those compounds in Table H which were not solid at room temperature are presented below.

5 Compound 907

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 2.4 (6H, s), 7.2-7.4 (4H, m), 7.95 (1H, s), 8.45 (1H, s).

Compound 908

<sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 2.3 (3H), 2.4 (3H), 7.1 (1H), 7.3 (3H, m), 7.8 (1H), 8.45 (1H).

10

Compound 909

*m/z* (EI) 382 (M<sup>+</sup>).

Compound 910

15 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 2.5 (3H), 7.45 (1H, d), 7.5-7.7 (2H, m), 7.75 (1H, d), 8.0 (1H, d), 8.5 (1H, d).

Compound 911

20 <sup>1</sup>H N.M.R (CDCl<sub>3</sub>) δ (ppm) 0.8 (t), 1.15 (d), 1.4 (quintet), 2.0 (s), 2.3 (s), 3.65 (dd), 7.7 (m), 7.95 (m).

Compound 912

*m/z* (EI) 357 ( $M^+$ ).

Compound 913

5 *m/z* (EI) 320 ( $M^+$ ).

Compound 914

*m/z* (EI) 330 ( $M^+$ ).

10 Compound 915

*m/z* (EI) 342 ( $M^+$ ).

Compound 916

*m/z* (EI) 315 ( $M^+$ ).

15

Compound 917

*m/z* (EI) 364 ( $M^+$ ).

Compound 918

20 *m/z* (EI) 364 ( $M^+$ ).

Compound 919

*m/z* (EI) 344 ( $M^+$ ).

25 Compound 920

$^1\text{H}$  N.M.R ( $\text{CDCl}_3$ )  $\delta$  (ppm) 2.35 (s), 2.5 (s), 7.4 (d), 7.8 (m), 7.9 (d).

Compound 921

30  $^1\text{H}$  N.M.R ( $\text{CDCl}_3$ )  $\delta$  (ppm) 3.9 (3H, m), 6.9-7.05 (2H, m), 7.5-7.75 (2H, m), 7.95 (1H, d),  
8.0 (1H, d), 8.5 (1H), 9.1 (1H).

Compound 922

*m/z* (EI) 328 ( $M^+$ ).

Compound 924

$m/z$  (EI) 382 ( $M^+$ ).

5 Compound 925

$^1H$  N.M.R (CDCl<sub>3</sub>)  $\delta$  (ppm) 1.3 (m), 2.7 (m), 4.2 (m), 7.75 (d), 7.95 (d).

Compound 927

$m/z$  (EI) 314 ( $M^+$ ).

10

Compound 928

$m/z$  (EI) 345 ( $M^+$ ).

Compound 93015 

$^1H$  N.M.R (CDCl<sub>3</sub>)  $\delta$  (ppm) 2.35 (d), 7.25 (m), 7.5 (d), 7.9 (d), 8.5 (d), 8.65 (s).

Compound 931

$m/z$  (EI) 301 ( $M^+$ ).

20 Compound 942

$^1H$  N.M.R (CDCl<sub>3</sub>)  $\delta$  (ppm) 2.6 (3H, s), 3.05 (3H, s), 8.0 (5H, m), 8.5 (1H, s).

Compound 943

$^1H$  N.M.R (CDCl<sub>3</sub>)  $\delta$  (ppm) 7.4-7.6 (7H, m), 7.8-8.0 (6H, m), 8.5 (1H, d).

25

Compound 944

$m/z$  (EI) 393 ( $M^+$ ).

Compound 94530 

$^1H$  N.M.R (CDCl<sub>3</sub>)  $\delta$  (ppm) 2.7 (3H, s), 7.15 (1H, dd), 7.5-7.65 (3H, m), 7.95 (1H, d), 8.1-8.25 (2H, m), 8.5 (1H, d).

Compound 946

$m/z$  (EI) 396 ( $M^+$ ).

Compound 947

5  $m/z$  (EI) 368 ( $M^+$ ).

Compound 948

$m/z$  (EI) 376 ( $M^+$ ).

10 Compound 949

$^1\text{H N.M.R}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 0.1-1.5 (5H, m), 1.7-1.9 (5H, m), 2.6 (1H, t), 8.0 (1H, m), 8.55 (1H, m).

Compound 950

15  $m/z$  (EI) 406 ( $M^+$ ).

Compound 951

$m/z$  (EI) 332 ( $M^+$ ).

20 Compound 952

$m/z$  (EI) 349 ( $M^+$ ).

Compound 953

25  $^1\text{H N.M.R}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 1.4 (3H, t), 3.0 (2H, q), 7.2 (3H, t) isomer, 7.3 (1H, d), 7.55 (1H, dd), 8.0 (1H, d, m), 8.55 (1H, d, m).

Compound 954

$^1\text{H N.M.R}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 2.5 (3H, m), 7.2-7.4 (2H, m), 7.5 (1H, d), 7.9 (1H), 8.4 (1H).

30 Compound 955

$^1\text{H N.M.R}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm) 2.6 (s), 4.8 (s), 7.25 (t), 7.5 (dd), 7.9 (m), 8.05 (d), 8.15 (d), 8.5 (s), 8.8 (d).

Compound 956

$m/z$  (EI) 374 ( $M^+$ ).

5    Compound 957

$m/z$  (EI) 314 ( $M^+$ ).

Compound 958

$^1H$  N.M.R ( $CDCl_3$ )  $\delta$  (ppm) 7.35-7.6 (8H, m), 7.9 (1H), 8.5 (1H).

10

Example 13

*N*-(3-Chloro-5-trifluoromethyl-2-pyridyloxy)-1-naphthalenecarboxamide

(Compound 1012)

A mixture of 1-naphthoic acid (0.46 g) and carbonyldiimidazole (0.44 g) in tetrahydrofuran  
15 (40 ml) was stirred for 16 hours under a nitrogen atmosphere. The product from stage b)  
(0.57 g) was then added, and the mixture stirred for 5 days. The solution was poured into  
saturated brine solution and the organic portion extracted with ethyl acetate (x3), dried  
( $MgSO_4$ ), filtered and evaporated. The residue was purified by silica gel chromatography  
(ethyl acetate/petrol) and triturated (diisopropyl ether) to give the title product, m.p. 198-9  
20 °C.

Preparation of Starting Materials

a)    2-{[3-Chloro-5-(trifluoromethyl)-2-pyridyl]oxy}-1,3-isooindolinedione  
2,3-Dichloro-5-trifluoromethylpyridine (50.0 g) was added over 5 minutes to a  
25 stirred solution of *N*-hydroxyphthalimide (37.5 g) and triethylamine (25.8 g) in  
acetone (750 ml). The mixture was refluxed for 8 hours and allowed to stand at  
room temperature for 16 hours. The solution was filtered and the filtrate evaporated  
to yield a solid which was partitioned between ethyl acetate and sodium bicarbonate  
solution. The organic fraction was isolated and the aqueous material re-extracted  
30 using further portions of ethyl acetate. The combined organic extracts were washed  
with water, dried, filtered and evaporated to give the crude product. The residue was  
triturated with diisopropyl ether to furnish the title compound as a white solid.



b) O-[3-Chloro-5-(trifluoromethyl)-2-pyridyl]hydroxylamine

Hydrazine monohydrate (1.7 g) was added to a solution of the product from stage a) (11.3 g) in tetrahydrofuran (200 ml) and the mixture stirred for 16 hours. The mixture was then filtered and the residual solid washed with a small volume of tetrahydrofuran and ethyl acetate, then four times with a 0.02M solution of sodium hydroxide saturated with sodium chloride. The combined aqueous layers were extracted with dichloromethane (x2) and the combined organic extracts dried, filtered and evaporated to give the title compound.


10 Example 14
N-(3-Chloro-5-trifluoromethyl-2-pyridyloxy)-N-methyl-1-naphthalenecarboxamide  
(Compound 1017)

Iodomethane (0.82 g) was added to a stirred solution of the product from Example 13 (Compound 1012) (1.93 g) and potassium *tert*-butoxide (0.61 g) in tetrahydrofuran (50 ml).

15 The reaction mixture was stirred for 48 hours. The solvent was evaporated and the residue partitioned between ethyl acetate and saturated aqueous ammonium chloride. The aqueous layer was separated and extracted with 3 portions of ethyl acetate. The combined organic phases were dried, filtered and evaporated to give a residue which was purified by silica gel chromatography (ethyl acetate/petrol) to give the title compound,  $m/z$  (EI) 380 ( $M^+$ ).

20

The following compounds of formula Ir (see Table J), i.e. compounds of general formula I where  $A^1$  is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is  $-O-N(R^3)C(=O)-$ , may be prepared by methods analogous to those of Examples 13 and 14.




(Ir)

Table J

Cmp	R <sup>3</sup>	A <sup>2</sup>	m.p.(°C)
1001	H	5-Me-2-pyrazinyl	202-6

25

Cmp	R <sup>3</sup>	A <sup>2</sup>	m.p.(°C)
1002	H	4-tolyl	190-3
1003	H	2-Cl-4-CF <sub>3</sub> -pyrimidin-5-yl	204-5
1004	H	4-Cl-phenyl	191-3
1005	H	2-NO <sub>2</sub> -5-(2-Cl-4-CF <sub>3</sub> -phenoxy)-phenyl	168-70
1006	H	3,5-diMe-4-isoxazolyl	108-11
1007	H	2,4-diMe-5-thiazolyl	152-5
1008	H	4,6-diMeO-2-( $\alpha,\alpha$ -diMe-4-Cl-benzyl)-pyrimidin-5-yl	124-5
1009	H	5-(3,5-diCl-phenoxy)-2-furyl	120-2
1010	H	6-MeO-3-pyridyl	157-9
1011	H	2-naphthyl	180
1012	H	1-naphthyl	198-9
1013	H	2-Cl-phenyl	170
1014	H	3-quinoliny	238-9
1015	H		oil
1016	H	4-morpholinyl-3-NO <sub>2</sub> -phenyl	217-8
1017	Me	1-naphthyl	oil
1018	H	1-naphthyl	218-20
1019	H	2,6-diCl-phenyl	246-7

The mass spectral data of the compounds in Table J which were not solid at room temperature are presented below.

5 Compound 1015

*m/z* (EI) 412 (M<sup>+</sup>).

Compound 1017

*m/z* (EI) 380 (M<sup>+</sup>).


Example 152-Methyl-1,2,3,4-tetrahydro-1-naphthalenone O-1-[3-chloro-5-(trifluoromethyl)-2-pyridyl]oxime5 (Compound 1101)

The starting material (0.58 g) was dissolved in tetrahydrofuran (5 ml) and to this was added potassium *tert*-butoxide (0.42 g) dissolved in tetrahydrofuran (5 ml). The mixture was stirred overnight and a solution of 2,3-dichloro-5-trifluoromethyl pyridine (0.72 g) in tetrahydrofuran (2 ml) was added. The mixture was stirred for 48 hours at room  
10 temperature, then the solvent was evaporated. The residue was partitioned between ethyl acetate and water. The organic layer was isolated, dried, filtered and evaporated to yield the title product as a light yellow gum,  $m/z$  (EI) 354 ( $M^+$ ).

a) 2-Methyl-1,2,3,4-tetrahydro-1-naphthalenone oxime







15 To a solution of 2-methyl-1-tetralone (3.20 g) in methanol (5 ml) was added hydroxylamine hydrochloride (1.81 g) in methanol (15 ml) and triethylamine (2.63 g). The mixture was stirred at 65°C for 5 hours, allowed to cool and stand at room temperature for 16 hours. The solvent was evaporated and water added to the residue. The product was extracted with ethyl acetate (3 portions) and the combined  
20 extracts were dried, filtered and evaporated to give an orange oil. On standing this separated into two layers. The top layer was removed and the bottom layer slowly solidified to give the title product as an orange solid.





The following compounds of formula Iq (see Table K), i.e. compounds of general formula I  
25 where  $A^1$  is 3-Cl-5-CF<sub>3</sub>-2-pyridyl and L is -O-N=C(R<sup>1</sup>)-, wherein R<sup>1</sup> and A<sup>2</sup>, together with the interconnecting atoms forms a 5- or 6- membered ring, may be prepared by methods analogous to those of Example 15.



(Iq)

Table K

Cmp	RZ	m.p.(°C)
1101	 <chem>CN=C1CCCc2ccccc12</chem>	oil
1102	 <chem>CN=C1CCCc2cc(OC)ccc12</chem>	oil
1103	 <chem>CN=C1CCCc2cc([N+](=O)[O-])ccc12</chem>	oil
1104	 <chem>CN=C1CCOCc2ccccc12</chem>	oil
1105	 <chem>CN=C1CCOCc2cc(Cl)ccc12</chem>	oil
1106	 <chem>CN=C1CCSCc2ccccc12</chem>	oil

Cmp	RZ	m.p.(°C)
1107		oil
1108		oil
1109		oil
1110		oil

Those compounds in Table K which do not have discrete melting points have the following characteristic mass spectral data.

5 Compound 1101

*m/z* (EI) 354 ( $M^+$ ).

Compound 1102

*m/z* (EI) 370 ( $M^+$ ).

10

Compound 1103

*m/z* (EI) 385 ( $M^+$ ).

Compound 1104

*m/z* (EI) 342 ( $M^+$ ).

Compound 1105

5 *m/z* (EI) 376 ( $M^+$ ).

Compound 1106

*m/z* (EI) 358 ( $M^+$ ).

10 Compound 1107

*m/z* (EI) 346 ( $M^+$ ).

Compound 1108

*m/z* (EI) 370 ( $M^+$ ).

15

Compound 1109

*m/z* (EI) 355 ( $M^+$ ).

Compound 1110

20 *m/z* (EI) 389 ( $M^+$ ).

Example 16

2-{{2-(3-Bromo-4-methoxyphenyl)-1*H*-1-imidazolyl}methyl}-3-chloro-5-  
(trifluoromethyl)pyridine

25 (Compound 1201)

To a solution of 2-(3-bromo-4-methoxyphenyl)-1*H*-imidazole (0.5 g) in tetrahydrofuran was added sodium hydride (0.08 g). After 30 minutes 3-chloro-2-(chloromethyl)-5-trifluoromethyl pyridine (0.46 g) was added and the solution heated until the reaction was complete. The reaction mixture was cooled, poured onto water and the organic phase  
30 extracted using dichloromethane, dried and evaporated to yield the crude product as an orange gum. Silica gel column chromatography yielded a gum which was further treated with diisopropyl ether and filtered. Evaporation of the filtrate afforded the title compound,  
*m/z* (APCI) 445 ( $M^-$ ).

WO 01/12604

2-(3-Bromo-4-methoxyphenyl)imidazole was synthesised from 3-bromo-4-methoxybenzonitrile using a method known to the skilled chemist.

### Test Example

5 Compounds were assessed for activity against one or more of the following:

*Phytophthora infestans*: late tomato blight

*Plasmopara viticola*: vine downy mildew

*Erysiphe graminis f. sp. tritici*: wheat powdery mildew

*Pyricularia oryzae*: rice blast

10 *Leptosphaeria nodorum*: glume blotch

Aqueous solutions or dispersions of the compounds at the desired concentration, including a wetting agent, were applied by spray or by drenching the stem base of the test plants, as appropriate. After a given time, plants or plant parts were inoculated with appropriate test pathogens before or after application of the compounds as appropriate, and kept under controlled environmental conditions suitable for maintaining plant growth and development of the disease. After an appropriate time, the degree of infection of the affected part of the plant was visually estimated. Compounds are assessed on a score of 1 to 3 where 1 is little or no control, 2 is moderate control and 3 is good to total control. At a concentration of 500 ppm (w/v) or less, the following compounds scored 2 or more against the fungi specified.

20 *Phytophthora infestans*: 49, 102, 119, 126, 202, 214, 215, 601, 902, 912, 927, 953, 1101 and 1102.

*Plasmopara viticola*: 5-7, 9, 10, 12, 102, 109, 126, 214, 215, 601, 901, 907, 914, 915, 921, 926-30, 958, 1001 and 1013.


25 *Erysiphe graminis f. sp. tritici*: 501, 901, 906, 913-5, 923, 926-931, 933, 935, 936, 948-50, 952, 954, 1008, 1102, 1104, 1107 and 1108.

*Pyricularia oryzae*: 7, 9, 11, 17, 126, 901, 906, 907, 913, 922, 923, 926-31, 937, 938, 939 and 1001.

30 *Leptosphaeria nodorum*: 23, 51, 53, 126, 207, 208, 906, 923, 926, 929, 933, 1007 and 1109.

Claims

- 5 1. The use of a compound of general formula I or salts thereof as phytopathogenic fungicides



where

10 A<sup>1</sup> is 2-pyridyl or its *N*-oxide, each of which may be substituted by up to four groups at least one of which is haloalkyl;

A<sup>2</sup> is optionally substituted heterocyclcyl or optionally substituted carbocyclcyl;

L is a 3-atom linker selected from the list: -CH(R<sup>1</sup>)N(R<sup>3</sup>)CH(R<sup>2</sup>)-,

15 -N(R<sup>3</sup>)N(R<sup>4</sup>)C(=X)-, -C(=X)N(R<sup>3</sup>)CH(R<sup>1</sup>)-, -CH(R<sup>1</sup>)OC(=X)-,  
 -CH(R<sup>1</sup>)OCH(R<sup>2</sup>)-, -N(R<sup>3</sup>)C(=X)N(R<sup>4</sup>)-, -C(R<sup>1</sup>)=C(R<sup>2</sup>)C(=X)-,  
 -C(R<sup>1</sup>)=N-N(R<sup>3</sup>)-, -CH(R<sup>1</sup>)N=C(R<sup>2</sup>)-, -O-N=C(R<sup>1</sup>)-, -O-N(R<sup>3</sup>)C(=X)-,  
 -N(R<sup>3</sup>)N(R<sup>4</sup>)CH(R<sup>1</sup>), -N(R<sup>3</sup>)C(Y)=N-, -N=C(Y)-N(R<sup>3</sup>)-, -N(R<sup>3</sup>)N=C(Y)-,  
 -C(=X)-N(R<sup>3</sup>)N(R<sup>4</sup>)-, -C(Y)=N-N(R<sup>4</sup>)- and -N(R<sup>3</sup>)CH(R<sup>1</sup>)C(=X)-;

wherein A<sup>1</sup> is attached to the left hand side of linker L;

20 where R<sup>1</sup> and R<sup>2</sup>, which may be the same or different, are R<sup>b</sup>, cyano, nitro, halogen, -OR<sup>b</sup>, -SR<sup>b</sup> or optionally substituted amino;

R<sup>3</sup> and R<sup>4</sup>, which may be the same or different, are R<sup>b</sup>, cyano or nitro;

or any R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> group, together with the interconnecting atoms, can form a 5- or 6-membered ring with any other R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup>, or any R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> or R<sup>4</sup> group, together with the interconnecting atoms can form a 5- or 6-membered ring with A<sup>2</sup>;

X is oxygen, sulfur, N-OR<sup>b</sup>, N-R<sup>b</sup> or N-N(R<sup>b</sup>)<sub>2</sub>; and

Y is halogen, -OR<sup>b</sup>, -SR<sup>b</sup>, -N(R<sup>b</sup>)<sub>2</sub>, -NR<sup>b</sup>(OR<sup>b</sup>) or -NR<sup>b</sup>N(R<sup>b</sup>)<sub>2</sub>;



wherein  $R^b$  is alkyl, alkenyl, alkynyl, carbocyclyl or heterocyclyl, each of which may be substituted; or hydrogen or acyl, or two adjacent  $R^b$  groups together with the nitrogen atom to which they are attached may form a 5- or 6-membered ring.

5

2. A pesticidal composition comprising at least one compound as claimed in claim 1 in admixture with an agriculturally acceptable diluent or carrier.

3. A method of combating pests at a locus infested or liable to be infested therewith,  
10 which comprises applying to the locus a compound as claimed in claim 1.

---

# INTERNATIONAL SEARCH REPORT

Intern. Application No  
PCT/EP 00/08268

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C07D213/61 C07D213/89 C07D405/12 C07D213/77 C07D401/12 C07D213/81 C07D213/64 C07D409/12 C07D417/12 C07D498/04 C07D401/06 A01N43/40		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, PAJ, CHEM ABS Data, WPI Data, BIOSIS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 99 42447 A (MOLONEY BRIAN ANTHONY ;SAVILLE STONES ELIZABETH ANNE (GB); AGREVO) 26 August 1999 (1999-08-26) the whole document;tautomers with Rb=H ----	1-3
X	EP 0 882 717 A (KYOWA HAKKO KOGYO KK) 9 December 1998 (1998-12-09) example 275 ----	1
X	EP 0 648 752 A (IHARA CHEMICAL IND CO ;KUMIAI CHEMICAL INDUSTRY CO (JP)) 19 April 1995 (1995-04-19) examples 302-307,345,346 ----	1-3
X	EP 0 573 883 A (BAYER AG) 15 December 1993 (1993-12-15) examples 172,173,245,246 ----- -/--	1-3
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *Z* document member of the same patent family		
Date of the actual completion of the international search  22 November 2000		Date of mailing of the international search report  05/12/2000
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer  Frelon, D

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/08268

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 469 711 A (SUMITOMO CHEMICAL CO) 5 February 1992 (1992-02-05) examples 90-92, 151 ---	1-3
X	EP 0 288 976 A (CIBA GEIGY AG) 2 November 1988 (1988-11-02) page 8 -page 15; examples ---	1-3
X	EP 0 270 061 A (HOFFMANN LA ROCHE) 8 June 1988 (1988-06-08) example 16 ---	1-3
X	DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MINN, KLEMENS: "Chalcones via a palladium-catalyzed coupling of iodoheterocycles to 1-phenyl-2-propyn-1-ol" retrieved from STN Database accession no. 115:8710 CA XP002152591 RN 134182-89-1 & SYNLETT (1991), (2), 115-16 , 1991, ---	1
X	PATENT ABSTRACTS OF JAPAN vol. 1995, no. 04, 31 May 1995 (1995-05-31) -& JP 07 025853 A (ISHIHARA SANGYO KAISHA LTD), 27 January 1995 (1995-01-27) example 21 ---	1-3
Y	WO 99 07687 A (AGREVO UK LTD ;COOPER IAN PAUL (GB); WEST PETER JOHN (GB); CARVER) 18 February 1999 (1999-02-18) example 19B; table B ---	1-3
Y	WO 98 50352 A (BRIGGS GEOFFREY GOWER ;CORNELL CLIVE LEONARD (GB); AGREVO UK LTD ( ) 12 November 1998 (1998-11-12) page 27; example 313 ---	1-3
Y	WO 98 42671 A (HAMPRECHT GERHARD ;BASF AG (DE); MENGES MARKUS (DE); WALTER HELMUT) 1 October 1998 (1998-10-01) the whole document ---	1-3
Y	WO 97 10215 A (BASF AG ;WAGNER OLIVER (DE); WETTERICH FRANK (DE); EICKEN KARL (DE) 20 March 1997 (1997-03-20) page 32 -page 33; table 4 ---	1-3
	-/--	

# INTERNATIONAL SEARCH REPORT

Intern. Application No  
PCT/EP 00/08268

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 92 07848 A (UNIROYAL CHEM CO INC ;UNIROYAL CHEMICAL LTD (CA)) 14 May 1992 (1992-05-14) abstract; claims ---	1-3
Y	EP 0 648 729 A (SUMITOMO CHEMICAL CO) 19 April 1995 (1995-04-19) pages 46, 49-77 ---	1-3
Y	EP 0 577 555 A (CIBA GEIGY AG) 5 January 1994 (1994-01-05) page 24; example 5.22 ---	1-3
Y	EP 0 350 691 A (BASF AG) 17 January 1990 (1990-01-17) page 8 -page 10 ---	1-3
A	EP 0 287 691 A (DOW CHEMICAL CO) 26 October 1988 (1988-10-26) example 14 ---	1-3
A	GB 2 307 177 A (AGREVO UK LTD) 21 May 1997 (1997-05-21) example 524 ---	1-3
A	GB 2 068 365 A (DOW CHEMICAL CO) 12 August 1981 (1981-08-12) the whole document ---	
A	PATENT ABSTRACTS OF JAPAN vol. 016, no. 148 (C-0928), 13 April 1992 (1992-04-13) -& JP 04 005282 A (MITSUBISHI PETROCHEM CO LTD), 9 January 1992 (1992-01-09) example 27 ---	1-3
A	PATENT ABSTRACTS OF JAPAN vol. 014, no. 310 (C-0736), 4 July 1990 (1990-07-04) -& JP 02 104575 A (ISHIHARA SANGYO KAISHA LTD), 17 April 1990 (1990-04-17) example 4 ---	1-3
A	PATENT ABSTRACTS OF JAPAN vol. 013, no. 379 (C-628), 22 August 1989 (1989-08-22) -& JP 01 131146 A (MITSUI PETROCHEM IND LTD;OTHERS: 01), 24 May 1989 (1989-05-24) example 21 ---	1-3
	-/--	

# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/EP 00/08268

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PATENT ABSTRACTS OF JAPAN  vol. 007, no. 114 (C-166),  18 May 1983 (1983-05-18)  -&amp; JP 58 035174 A (ISHIHARA SANGYO KK),  1 March 1983 (1983-03-01)  abstract</p> <p>-----</p>	1-3

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Present claims 1-3 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely the examples and closely related homologues wherein A1 represents 3-chloro-5-trifluoro-pyrid-2-yl and A2 is (opt. substit.) phenyl, pyridyl, pyrimidyl, pyrazinyl, furanyl, thienyl, (iso)thiazolyl, (iso)oxazolyl.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/EP 00/08268

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9942447 A	26-08-1999	AU 2527199 A NO 20004159 A	06-09-1999 17-10-2000
EP 0882717 A	09-12-1998	AU 719392 B AU 4470897 A NZ 330571 A CA 2239227 A CN 1208404 A WO 9814431 A	11-05-2000 24-04-1998 28-10-1999 09-04-1998 17-02-1999 09-04-1998
EP 0648752 A	19-04-1995	BR 9405139 A DE 69421036 D DE 69421036 T RU 2131421 C US 5616594 A CN 1105179 A ES 2139071 T WO 9424110 A JP 6345608 A US 5700824 A	31-08-1999 11-11-1999 08-06-2000 10-06-1999 01-04-1997 12-07-1995 01-02-2000 27-10-1994 20-12-1994 23-12-1997
EP 0573883 A	15-12-1993	DE 4219247 A AU 3877693 A JP 6065219 A	16-12-1993 16-12-1993 08-03-1994
EP 0469711 A	05-02-1992	AU 638840 B AU 7913691 A BR 9102835 A CA 2046206 A JP 5004971 A US 5135563 A	08-07-1993 09-01-1992 04-02-1992 06-01-1992 14-01-1993 04-08-1992
EP 0288976 A	02-11-1988	AT 106394 T AU 616176 B AU 1525088 A BR 8802063 A DE 3889767 D ES 2053612 T IL 86189 A JP 2575046 B JP 63284162 A KR 9700951 B PT 87341 A, B US 5015649 A US 5126358 A ZA 8803029 A	15-06-1994 24-10-1991 03-11-1988 29-11-1988 07-07-1994 01-08-1994 15-07-1992 22-01-1997 21-11-1988 21-01-1997 01-05-1988 14-05-1991 30-06-1992 01-11-1988
EP 0270061 A	08-06-1988	AU 602372 B AU 8203687 A DK 570987 A HU 45850 A US 4943583 A BR 8706464 A JP 63154678 A ZA 8708811 A	11-10-1990 02-06-1988 02-06-1988 28-09-1988 24-07-1990 12-07-1988 27-06-1988 01-06-1988
JP 07025853 A	27-01-1995	NONE	

# INTERNATIONAL SEARCH REPORT

information on patent family members

Intern. Application No

PCT/EP 00/08268

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9907687 A	18-02-1999	AU 8635298 A	01-03-1999
WO 9850352 A	12-11-1998	AU 7220498 A	27-11-1998
WO 9842671 A	01-10-1998	AU 6829298 A	20-10-1998
		EP 0968188 A	05-01-2000
WO 9710215 A	20-03-1997	DE 19533653 A	13-03-1997
		DE 19535208 A	27-03-1997
		AU 6987096 A	01-04-1997
		EP 0852579 A	15-07-1998
		JP 11512406 T	26-10-1999
		US 5955473 A	21-09-1999
WO 9207848 A	14-05-1992	US 5071862 A	10-12-1991
		AT 115578 T	15-12-1994
		AU 9017891 A	26-05-1992
		CN 1062907 A	22-07-1992
		DE 69105995 D	26-01-1995
		DE 69105995 T	11-05-1995
		DK 555365 T	15-05-1995
		EP 0555365 A	18-08-1993
		IL 99868 A	08-12-1995
		MX 9101775 A	05-06-1992
		ZA 9108572 A	26-08-1992
EP 0648729 A	19-04-1995	AU 675580 B	06-02-1997
		AU 7588094 A	11-05-1995
		BR 9404147 A	13-06-1995
		CA 2118349 A	20-04-1995
		CN 1108642 A	20-09-1995
		DE 69411658 D	20-08-1998
		DE 69411658 T	25-02-1999
		EG 20417 A	28-02-1999
		ES 2119046 T	01-10-1998
		JP 7188088 A	25-07-1995
		RU 2130008 C	10-05-1999
		TR 28760 A	28-02-1997
		US 5530015 A	25-06-1996
		US 5698702 A	16-12-1997
		ZA 9408162 A	19-06-1995
EP 0577555 A	05-01-1994	AT 159930 T	15-11-1997
		AU 4160593 A	06-01-1994
		BR 9302700 A	02-08-1994
		CA 2099309 A	31-12-1993
		CN 1083804 A, B	16-03-1994
		DE 59307623 D	11-12-1997
		DK 577555 T	27-07-1998
		ES 2110587 T	16-02-1998
		GR 3025440 T	27-02-1998
		HR 931007 A	29-02-1996
		HU 65419 A	28-06-1994
		JP 2855398 B	10-02-1999
		JP 6073036 A	15-03-1994
		MX 9303929 A	31-01-1994
		NZ 248008 A	26-01-1996
		SG 49660 A	25-05-1999



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/EP 00/08268

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0577555 A		TR 26865 A	19-08-1994
		US 5481013 A	02-01-1996
		ZA 9304641 A	05-01-1994
EP 0350691 A	17-01-1990	DE 3823991 A	15-02-1990
		AT 123023 T	15-06-1995
		AU 616031 B	17-10-1991
		AU 3811689 A	01-02-1990
		DE 58909248 D	29-06-1995
		ES 2072276 T	16-07-1995
		JP 2088558 A	28-03-1990
		JP 2871732 B	17-03-1999
		MX 9203412 A	31-08-1992
		NZ 229933 A	26-02-1991
		US 5157037 A	20-10-1992
		US 5194438 A	16-03-1993
		US 5326767 A	05-07-1994
		US 5378711 A	03-01-1995
EP 0287691 A	26-10-1988	CA 1303617 A	16-06-1992
		US 4670045 A	02-06-1987
GB 2307177 A	21-05-1997	NONE	
GB 2068365 A	12-08-1981	US 4331669 A	25-05-1982
		US 4472583 A	18-09-1984
JP 04005282 A	09-01-1992	NONE	
JP 02104575 A	17-04-1990	NONE	
JP 01131146 A	24-05-1989	NONE	
JP 58035174 A	01-03-1983	NONE	